

Microbial quality of catfish (*Clarias Gariepinus*) smoked with Nigerian stored products research institute (Nspri) developed smoking kiln

October 17, 2013 Volume 2; Issue 3

Adedayo M Rachael, Olayemi F Foline, Bamishaiye E Iyabo, Ajiboye A Elizabeth

Rachael AM, Foline OF, Iyabo BE, Elizabeth AA. Microbial quality of catfish (*Clarias Gariepinus*) smoked with Nigerian stored products research institute (Nspri) developed smoking kiln. Bioresearch Bulletin. 2013 Oct 17 [last modified: 2013 Oct 17]. Edition 1.

Abstract

The microbial quality of cat fish (*Clarias gariepinus*) smoked with Nigerian Stored Products Research Institute (NSPRI) developed smoking kiln was assessed. Raw fish samples were purchased and cleaned before smoking in the kiln designed and constructed by the institute. Potato dextrose agar and nutrient agar were used for the mould/yeast and bacteria load of the smoked fish respectively while appropriate selective media were used for the isolation of the enteric organisms. The total bacteria count was 2.0×10^{-4} cfug and 0.7×10^{-4} cfug for mould/yeast. The result showed that no *Pseudomonas*, *Escherichia coli* or *Salmonella* species was found on the smoked fish. Generally the total bacteria count was higher than the mould/yeast count. The sensory evaluation of the smoked fish showed that the fish was generally acceptable in taste and flavour. In conclusion, *C. gariepinus* smoked with NSPRI developed smoking kiln is of high microbiological standard and suitable for human and livestock consumption.

Introduction

Cat fish (

Clarias gariepinus

) is a very important freshwater fish in Nigeria. It has enjoyed wide acceptability in most parts of the country because of its unique taste, flavour and good texture. It is widely distributed, extensively cultivated in ponds. Fish is one of the best source of proteins, vitamins and minerals and are essential nutrients required for supplementing both infant and adult diets¹. In Nigeria, fish is eaten fresh, preserved or processed (smoked) and form a much-cherished delicacy that cuts across socio-economic, age, religious and educational barriers². The microbial flora associated with freshly harvested fish is principally a function of the environment in which the fish are caught and not of the fish species; hence, the indigenous microbial populations of fish can vary significantly³.

Fish, because of their soft tissues and aquatic environment are extremely susceptible to microbial contamination. Millions of bacteria, many of them potential spoilers, are present in the surface slime, on the gills and in the intestines of live fish, although the flesh itself is normally sterile. Bacterial growth and invasion on the fish are prevented by the body's natural defense system during life but after death the defense system breaks down and the bacteria multiply and invade the flesh⁴. Poor post-harvest technology (handling, preservation and processing) have ability to cause unhealthy situation through massive spoilage. An estimate of 40% postharvest losses of total fish landings have been reported in Nigeria⁵. Mayboom⁶ similarly reported that

15% of the total fish catch in Kainji Lake is lost because of spoilage and breakage between the sources of supply and the consumers. Saliu⁷, also reported that fish spoilage in Nigeria is influenced to a large extent by high ambient temperatures, considerable distances of landing ports to points of utilization and poor as well as inadequate infrastructure for post-harvest processing and landing. Bacterial spoilage is characterized by softening of the muscle tissue and the production of slime and offensive odors⁸. Among the several methods of long term preservation of fish, smoking is perhaps the simplest method as it does not require sophisticated equipment or highly skilled workers⁹. Smoking is a century old method of food preservation. Fish smoking is one of the traditional fish processing methods aimed at preventing or reducing postharvest losses. Smoking involves heat application to remove water and it inhibits bacterial and enzymatic actions of fish¹⁰¹¹¹²¹³¹⁴¹⁵¹⁶¹⁷ also noted that apart from giving the product a desirable taste and odor, smoking provides a longer shelf-life through its anti-bacterial and oxidative effect, lowering of pH, imparting desirable colorations as well as accelerating the drying process and acting as antagonist to spoilage. However, smoking is the most popular method of fish processing¹⁶. Traditionally, fish is smoked in pits or on raised smoking “tables” where the control of heat is difficult and at times impossible¹⁸. Also, the smoked fish is often exposed to dust and heavy microbial contamination during the process of traditional smoking which gives the final products high microbial load even immediately after smoking, this is in agreement with the findings of¹⁹ who reported that lack of proper smoking and proper hygienic handling of smoked fish products would result in a very high microbial load and open flame smoking of fish has been noted to produce cancer promoting compounds in the body. In an attempt to reduce the problems often encountered during smoking, Nigeria Stored Products Research Institute (NSPRI) fish smoking kiln was developed. Due to handling and the exposure to the environment during and after smoking, smoked fish often comes out with heavy microbial load after smoking therefore. This research is thus aimed at studying the microbial quality of

C. gariepinus

smoked in NSPRI designed and constructed smoking kiln.

Materials and Methods

Description of NSPRI kiln

The smoking kiln is rectangular in shape of dimension 60 x 60 x 120 cm in depth, width and height respectively. The structure has an internal wall made of galvanized iron (GI) sheet, lagged with 2.54 cm asbestos particles and covered with ½ in plywood. It has a unit compartment for drying, heating and smoking with removable four trays made from 25.81 cm square wire mesh placed on 2.54 cm angle iron. The structure was incorporated with three axial fan powered by ten 1.5 V battery to increase air supply to the heating chamber. The heating chamber was also made up of two sections which may be used together or independently, they are the heat producing section which can produce heat using charcoal, coal or coconut shell; and the smoke producing section using sawdust. Four of the ½ in diameter pipe exits were also provided at the top of the structure to aid the escape of water vapour during the smoking process while 6 of the ½ in diameter pipe were provided for air inlets. The structure works by burning charcoal to produce heat for the drying of the fish and by burning sawdust to generate smoke which imparts desired aroma and color to the smoked fish. The structure may be used to dry the fish alone without smoking and vice versa.

Fish processing

Sixty seven freshly harvested Catfish (

Clarias gariepinus

) aged six months were obtained from Fagam Farm through Kano State Department of Fisheries in Kano State, Nigeria. All were killed, beheaded, eviscerated and washed thoroughly. They were brined using a teaspoon of salt per fish. They were packaged into NSPRI Fish boxes which were insulated and the fish to ice in ratio 1:1 and

were transported to the laboratory for further treatments. The fish smoking kiln was operated by first loading charcoal into the heat chamber, preheating for some minutes, and then loading the salted pieces of fish onto the trays in its central chamber. The kiln was closed for some time to allow the smoking to take place. The capacity was 100 kg per barge. The smoking time, temperature and ambient conditions were monitored during the smoking operations.

The smoking was terminated when the fish were properly dried after 15 h. The fresh and smoked samples were analyzed for total microbial load, presence of yeast/mould and indicator organisms like

E. coli

and

Salmonella

species. The sensory evaluation was also conducted using nine unit hedonic scales conducted by six semi trained panels.

Microbial analysis

One gram (1g) representative sample was obtained aseptically from the muscle of the smoked catfish samples. The samples were grounded and serial dilutions (10^{-1} - 10^{-4}) of the homogenized samples were made using sterile distilled water. All chemicals used were of analytical grade and supplied by Sigma Co. (St Louis, USA). Each analysis was carried out in triplicates. All microbial analysis was done following the methods prescribed by 20.

Total Plate Count (TBC)

This was done using the pour plate method of 20. One milliliter of the serially diluted samples was taken in duplicates and plate count agar was poured at 40°C on the plates. The samples and the medium were properly mixed, allowed to set and incubated at 35°C for 24h. The number of colonies on the plates was counted.

Escherichia coli count

This was done using Eosin Methylene Blue Agar at 35°C for 24h. Colonies with green metallic sheen were counted as *E. coli*

Salmonella count

Samples for detection of

salmonella

were plated out on Salmonella-Shigella Agar. The plates were incubated at 35°C for 24h. Black colonies showed the presence of

Salmonella

sp.

Pseudomonas count

This was done using

Pseudomonas

(Specific Agar PSA). Colonies with blue-green or greenish yellow appearance which fluorescence were counted as

Pseudomonas.

Yeast and mould Counts.

This was done by plating out serially diluted samples on Potato Dextrose Agar (PDA) at room temperature (30-35°C) for 48-72h.

Discussion

The design of the NSPRI constructed smoking kiln ensures that the fish was properly smoked within 15 hours at 70°C. The moisture content of the fish was drastically reduced as shown by the reduction in weight and moisture content before and after smoking. Since dryness of 7.3% was obtained from 78.7% initial moisture content after 15 hours of drying, it implies that the drying rate of the catfish using the kiln was 4.76% moisture per hour. The difference in moisture content after smoking is a confirmation of the fact that

C. gariepinus

has high water content which predisposes it to high microbial spoilage if not well preserved after harvest. The moisture content of the smoked fish which was 7.3% falls within the allowable limit (6-8%) for smoked dried fish as this is of paramount importance in preventing spoilage during storage and enhances shelf life of the smoked fish, this observation is in agreement with the findings of²¹ and¹⁰ who reported that spoilage of fish resulting from the action of enzymes and bacteria can be slowed down by the addition of salt as well as reduction in moisture through sun drying or smoking. The total mean bacteria count was 2.0×10^4 colony forming unit per gramme of the fish sample, this value falls within the maximum recommended value of bacteria count for good quality fish products which is 5×10^{-5} colony forming unit per gramme according to²² and the Microbiological Guideline for Ready to eat - Food which is $< 10^6$ ²³. The result also indicated that there was no contamination with enteric organisms by handlers during smoking as there was no coliform found after smoking. The absence of

E. coli

and

Salmonella

species which are indicative organism of contamination by microorganisms from enteric origin further confirms the effectiveness of the smoking kiln. This was achieved as a result of the fact that fish smokers are not in direct contact with the fish during smoking. According to¹⁴ microbial action plays a large part in the spoilage of fish and fish products. Fish smoking in NSPRI developed kiln has been able to effectively reduce this main source of spoilage. It was also noticed that

Pseudomonas,

an opportunistic bacteria in food spoilage and infection was not found²⁴. The value of yeast/ mould recorded after smoking was also found to be within the acceptable limit²²²³. The sensory evaluation as shown in Table 3 which was conducted by six semi-trained panelist on a 9 point hedonic scale of smell, texture, colour taste and general acceptance showed that, the fish were in good quality and condition and can be widely accepted.

Conclusion

Considering the result above the NSPRI developed smoking kiln has proof to be effective in smoking fish and reducing its original microbial load to microbiologically recommended safe level thus making it fit for immediate consumption or storage for long time use or for export.

References

1. Abdullahi SA, Abolude DS, Ega RA. 2001. Nutrient Quality of four Oven Dried Freshwater Catfish in Northern Nigeria. J. Trop. Biosci., 70.
2. Adebayo-Tayo BC, Onilude AA, Patrick UG. 2008. Mycoflora of Smoke-dried Fishes Sold in Uyo, Eastern Nigeria. World J. Agric Sci.,

3. Shewan JM. 1961. The Microbiology of Sea Water Fish. In Fish as Food. Ed by G.Borgstrom, New York Academic press. 487.
 4. Abolagba OJ, Uwagbai EC. 2011. A Comparative Analysis of the Microbial Load of smoke-Dried Fishes (*Ethmalosa fimbriata* and *Pseudotolithus elongatus*) Sold in Oba and Koko Markets in Edo and Delta States, Nigeria at Different Seasons. Australian Journal of Basic and Applied Sciences 5(5):544-550.
 5. Akande GR. 1996. "Post-harvest processing in Fisheries". A paper presented at training for officers of UNDP assisted programme on artisanal fisheries development, Ogun State at Federal College of Fisheries and Marine Technology, Lagos. Jan. 15th - Feb. 5th, 1-20.
 6. Mayboom J. 1974. Fish handling and processing in the Kainji Lake Basin and suggestions for improvement and future research. Kainji Lake Research Institute, New Bussa, Nigeria. 34-45.
 7. Saliu JK. 2008. Effect of smoking and frozen storage on the nutrient composition of some African fish. Adv. Nat. Appl. Sci., 2(1):16-20.
 8. Geoff A, Ivors SS, Paul. 1991. Post-Harvest Losses of Fish in the tropics. Natural Resources Institute. 1-5
 9. Olayemi FF, Adedayo MR, Bamishaiye EI, Awagu EF. 2011. Proximate composition of catfish (*Clarias gariepinus*) smoked in Nigerian stored products research institute (NSPRI): Developed kiln. International Journal of Fisheries and Aquaculture 3(5):95-97.
- REFERENCE LINK
10. Kumolu-Johnson CA, Aladetohun NF, Ndimele PE. 2009. The effects of smoking on the Nutritional Qualities and Shelf-Life of *Clarias gariepinus* (BURCHELL 1822). African Journal of Biotechnology 9(1):73-76.
 11. Abolagba OJ, Melle OO. 2008. Chemical composition and keeping qualities of a scaly fish tilapia, *Oreochromis niloticus* smoked with two energy sources. Afr. J. Gen. Agric. 4(2):113-117.
 12. Olorok JO, Ihuahi JA, Omojowo FS, Falayi BA, Adelowo EA. 2007. Handbook of Practical Fisheries Technology. Published by Fisheries Technology Division, National Institute for Freshwater Fisheries Research (NIFFR), P.M.B 6006, New Bussa, Niger State. 22-29.
 13. Sengor GF, Kalafatoglu H, Gun H. 2004. The determination of Microbial flora, water activity and chemical analysis in smoked Mussels (*Mytilus galloprovincialis*, L.) Turk. J. Vit. Anim. Sci., (28):793-797.
 14. Eyo AA. 2001. Fish processing Technology in the Tropics. Published by National Institute for Freshwater Fisheries Research (NIFFR), P.M.B. 6006, New Bussa, Niger State.
 15. Horner WFA. 1997. Preservation of Fish by Curing (Drying, Salting and Smoking). In Fish processing Technology 2nd ed. Edited by G.M. Hall. 34-36.
 16. Olley J, Doe PE, Heruwati ES. 1988. The influence of drying and smoking on the nutritional properties of fish. In Burth JR (ed); An Introd. overview in fish smoking and drying. Elsevier, London. 1-14.
 17. Clucas IJ, Ward AR. 1996. Post Harvest Fisheries Development. A Guide to Handling, Preservation, Processing and Quality. Chatham Maritime, Kent ME4TB, United Kingdom. 665.
 18. Afolabi OA. 1984. Quality changes in Nigeria Traditionally Processed Freshwater species. J. Food Technol., 18:17-29.
 19. Abolagba OJ, Iyeru OA. 1998. Study of Insect Pest Infecting Traditionally Processed Fish Sold in Benin City Metropolis, Nigeria. Nig. J. Applied Sci., 16:25-29.
 20. Association of Official Analytical Chemists. 2000. Official Methods of Analysis of the Association of Official

Analytical Chemists, Vols. I & II, Association of Analytical Chemists, Arlington. 17th Edition.

21. Salán OE, Juliana AG, Marilia O. 2006. Use of smoking to add value to salmoned trout. Braz. Arch. Biol. Technol., 49(1):57-62.
22. International Commission on Microbiology Safety for Foods (ICMSF). 1986. Microorganisms in Food 2. Sampling and Specific Applications 2nd Edn. University, Toronto Press, Canada.
23. Microbiological Guideline for Ready to -eat - Food. 2007. Centre for Food Safety, Food and Environmental Hygiene Department. 43/F, Queensway Government Offices, 66 Queensway, Hong Kong.
24. Talaro KP. 2009. Foundations' in Microbiology, 7th Edition. McGraw-Hill Companies. Avenue of the Americas, New York. 595-596.