

Molecular pathobiology of Diabetic Retinopathy

November 23, 2010 · Volume 1; Issue 3

Ehtishamul Haq, Mujeeb Z Banday, Asma Bashir

Haq E, Banday MZ, Bashir A. Molecular pathobiology of Diabetic Retinopathy. Bioresearch Bulletin. 2010 Nov 23 [last modified: 2013 Oct 16]. Edition 1.

Abstract

Background: Diabetes and its associated complications constitute a huge global health menace. Hyperglycemia induces alterations in several signaling pathways, which ultimately culminate in the development of various complications that includes Diabetic Retinopathy (DR). DR, a disease of retina is a major cause of blindness among diabetics.

Challenges: Although the precise mechanism involved in the pathogenesis of DR is broadly unclear due to its multifactorial nature, the contribution of oxidative stress and inflammation occupy the central stage.

Conclusion: In this review, the prevalence, clinical and pathobiological characteristics of DR and the possible cross-talk between the different contributing pathways involved in its development are described.

Introduction

Diabetes mellitus or simply diabetes is a lifelong progressive disease that results from body's inability to produce insulin, use of insulin to its full potential or both, and is characterized by high circulating glucose (Hyperglycemia)³. This disease has reached epidemic proportion and has become one of the most challenging health problems of the 21st century. Indeed, by the year 2010, the total number of people with diabetes is projected to reach 221 million worldwide⁷ and by the year 2025, this number is estimated to reach 334 million. There are two distinct forms of diabetes, type 1, juvenile or Insulin-Dependent Diabetes Mellitus (IDDM) and type 2, adult onset or Non Insulin-Dependent Diabetes Mellitus (NIDDM)³. Type 2 diabetes accounts for 90 percent of all diabetes cases. Diabetes is a chronic heterogeneous metabolic disorder with complex pathogenesis. The sustained hyperglycemia in diabetes attacks both microvessels and macrovessels throughout the body that lead to complications, such as cardiovascular disease, renal failure, retinopathy and peripheral and autonomic neuropathy¹⁶³⁴.

As a consequence of these complications, diabetes is the leading cause of blindness and visual impairment, noninjury amputation, and end-stage kidney disease in adults at developed countries. The patients with diabetes develop cataracts in an earlier age, and are nearly twice as likely to get glaucoma compared to nondiabetics¹⁰⁰. These complications occur in both type 1 and type 2 diabetes irrespective of the fact that type 1 and type 2 diabetes originate from different pathogenetic causes thereby indicating that there is a significant association between hyperglycemia and the diabetic microvascular complications in both diabetes types¹².

Although, all the features of diabetic retinopathy may be found in both types of diabetes, but characteristically, the incidence of the main causes of vision loss, macular edema and retinal neovascularization is quite different for each type of diabetes⁶.

Diabetic Retinopathy

Diabetic retinopathy (DR), a disease of the retina, is the most common and specific microvascular complication of diabetes³³. It is the major cause of blindness among people in working age and is a leading cause of visual loss in older patients in developed countries¹⁰⁶, thus representing a major socioeconomic problem. Diabetic retinopathy is present to some degree in nearly all individuals who have had diabetes for more than 15 years, regardless of the type of diabetes. It is a duration-dependent disease that develops in stages; the incidence of retinopathy is rarely detected in the first few years of diabetes, but the incidence increases to 80% by 10 years, and to virtually 95% by 25 years of type 1 diabetes⁶. However, the overall incidence of DR is slightly less common in type 2 diabetes and nearly 75 per cent of type 2 diabetes patients develop this disease after 15 yr duration of diabetes⁵⁷, but it still remains the most frequent microvascular complication suffered by these patients. The pathogenic basis of diabetic retinopathy is not wholly understood at the cellular and molecular level but it is clear that the pathogenesis of diabetic retinopathy is highly complex and multifactorial. It is now known through major international epidemiological trials that hyperglycemia is a principle and underlying cause of DR in both IDDM and NIDDM². Hyperglycemia is responsible for not only microvascular¹² disease but also of macrovascular complications⁷¹ in both forms of diabetes. Hyperglycemia causes retinal damage via many acute (and repeated) and also cumulative long-term changes (Fig. 1).




Fig. 1: Hyperglycemia induced retinal damage via sustained acute and cumulative changes in cellular metabolism and subsequent cumulative changes in stable macromolecules that ultimately lead to tissue damage.

Diabetic retinopathy and vascular damage

In diabetic retinopathy, most cells of the retina are affected by the metabolic abnormalities of diabetes⁷⁶, but the sight-threatening manifestations of diabetic retinopathy are ultimately attributable to capillary damage. The blood vessels of retina have tight junctions that protect them from leaking, but sustained high glucose damages the tight junctions and the vessels become leaky that cause the blood vessels to swell and leak fluid. The damaged microvasculature allows the fluid or blood to seep into the retina resulting in the breakdown of blood–retinal barrier that causes the swelling of the retina or macular edema²⁴⁴⁴. The capillaries of retina are lined with endothelial cells that are responsible for maintaining the blood retinal barrier, and are supported with an equal number of pericytes (microvascular mural cells) that help provide tone to the vessels.

However, in diabetes, the ratio of endothelial cells to pericytes is altered to 4:1¹⁹⁵. As the disease progresses, the endothelial cells try to repair the damaged vessel by multiplying on the inner side of the vessel wall blocking the capillaries. Due to progressive dysfunction, the capillaries die prematurely leading to ischemia that is followed by new vessel growth or neovascularization. New capillaries start to grow from the surface of the retinal veins towards the center of the eye with no support, and ultimately lead to the detachment of the retina³⁷⁶³⁶¹⁰. The leukocytes become less deformable, and retinal leukostasis is increased affecting endothelial function⁵². The macular edema caused due to a breakdown in the blood–retinal barrier, is one of the earliest clinically detectable signs of DR and of all the clinical features, macular edema is most closely correlated with the degree of vision loss⁸⁴.

Classification of Diabetic Retinopathy

The diabetic retinopathy is broadly classified as either Non Proliferative Diabetic Retinopathy (NPDR) or Proliferative Diabetic Retinopathy (PDR). The classification of diabetic retinopathy has evolved as our understanding of diabetic eye disease has increased. Nonproliferative diabetic retinopathy, also known as background retinopathy, occurs when there are only intraretinal microvascular changes, such as altered retinal vascular permeability and eventual retinal vessel closure⁸⁰. It is further divided into NPDR with maculopathy, NPDR without maculopathy and pre-proliferative retinopathy. NPDR indicates progressive ischemia in the retina and an increased risk for the development of PDR and blindness.

The prominent clinical features of NPDR are often classified as red lesions, such as microaneurysms, dot or blot hemorrhages, and intraretinal microvascular abnormalities, or bright lesions, such as lipid or lipoprotein hard yellow exudates, and superficial retinal infarcts (cotton-wool spots)⁷⁵⁹⁰. Microaneurysms are saccular outpunching of the capillary wall in the region of vascular occlusions. These are visible as small round dark red dots on the retinal surface. Hemorrhages and exudates result from the damaged vasculature. Increase in retinal blood flow and vasodilation occurs early in the course of the disease, followed by the microvascular leakage and occlusion, and histologic data corroborates the clinical evidence³²⁷⁰. Maculopathy is defined as the presence of edema and/or haemorrhages and/or exudates and/or retinal thickening within 500 m (i.e., 1 disc diopter) of the fovea, with or without visual loss. Preproliferative diabetic retinopathy is the stage before the onset of neovascularization and is characterized by:

- (a) Extensive retinal haemorrhages,
- (b) Marked venous beading,
- (c) Numerous cotton wool spots or retinal infarcts,
- (d) Intra-Retinal Microvasculature Abnormalities [(IRMAs, a term chosen so as to be neutral about whether these abnormal vessels represent intraretinal new vessels or dilated preexisting vessels], and
- (e) Marked retinal ischemia as evidenced by capillary drop outs in the fundus fluorescein angiogram. Neovascularization is not a component of the nonproliferative phase.

However, in advanced NPDR, nonperfusion of the retina may develop and lead to the proliferative phase⁸⁰.

Proliferative diabetic retinopathy continues to be a major cause of blindness throughout the world. It is characterized by new vessels (neovascularization), preretinal haemorrhage, vitreous haemorrhage, vitreoretinal traction, localized retinal detachment and sometimes fibrous bands proliferating on the retinal surface and these are the primary complications leading to vision loss²⁶. In both nonproliferative and proliferative diabetic retinopathy, macular edema can occur as increased retinal vascular permeability leads to accumulation of fluid in the retinal area serving central vision⁸⁰. There is accumulated evidence indicating that only the Non Proliferative stage of Diabetic Retinopathy (NPDR) is directly due to the systemic disease and associated hyperglycemia and other metabolic alterations. Proliferative retinopathy occurs in diabetic eyes only after the development of widespread ischemia due to capillary closure. Neovessels in the retina are a direct result of retinal ischemia and not influenced by the diabetic metabolic control. Its course and management are not different from other situations in the retina where there is abnormal new vessel formation⁵⁴.

Oxidative stress in Diabetic Retinopathy

The pathogenic basis of diabetic retinopathy is not wholly understood but various studies have shown the important, in fact, the central role that oxidative stress and inflammation play in the pathogenesis of this disease. Oxidative stress refers to the condition in which there is a serious imbalance between the production of oxidants (reactive oxygen species (ROS) and reactive nitrogen species (RNS)), and antioxidant defense, leading to potential tissue damage⁹⁷. Diabetes results in increased oxidative stress and increased oxidative stress is one of the key regulators in the development of diabetic complications. The possible sources of oxidative stress in diabetes might include auto-oxidation of glucose, disruption of the mitochondrial electron transport chain, shifts in redox balances, decreased tissue concentrations of low molecular weight biologically existing antioxidants such as reduced glutathione (GSH), vitamin E, vitamin C and β -carotene and impaired activities of antioxidant defense enzymes such as superoxide dismutase (SOD), glutathione reductase, glutathione peroxidase and catalase, that are responsible for scavenging free radicals and maintaining redox homeostasis and as such superoxide levels are elevated in the mitochondria¹³ ¹³⁴⁵⁶⁰⁶¹⁶³⁶⁵⁶⁶⁸⁵⁸⁸⁵⁵. The retina has a high content of polyunsaturated fatty acids and has the highest oxygen uptake and glucose oxidation relative to any other tissue. This phenomenon renders retina more susceptible to oxidative stress⁸.

Hyperglycemia induced ROS generation is considered as a causal link between elevated glucose and the other metabolic abnormalities important in the development of diabetic complications and normalizing mitochondrial ROS is shown to prevent glucose induced metabolic abnormalities that are postulated in the pathogenesis of diabetic complications including the diabetic retinopathy⁸⁶¹⁹²⁰. The correlation between hyperglycemia, changes in the redox homeostasis, and oxidative stress have been suggested to be the key events in the pathogenesis of diabetic retinopathy. Animal studies have demonstrated that oxidative stress contributes not only to the development of diabetic retinopathy but also to the resistance of retinopathy to reverse after good glycemic control is reinstated—the metabolic memory phenomenon⁶².

Resistance of diabetic retinopathy to reverse is probably attributed to accumulation of damaged molecules and ROS that are not easily removed even after good glycemic control is reestablished. Since oxidative stress represents an imbalance between excess formation and/or impaired removal of ROS, the antioxidant defense system of the cell is a crucial part of the overall oxidative stress experienced by a cell. Further, the cell is equipped with intracellular antioxidant, GSH; GSH is probably the most important defense the cell is equipped with. It can act as an ROS scavenger and modulate intracellular redox state⁷⁸. The levels of this intracellular antioxidant are decreased in the retina in diabetes⁵⁶, and the enzymes responsible for its metabolism are compromised⁵⁹⁶⁴. Apart from the antioxidant defense enzymes, nonenzymic antioxidants such as vitamin C, vitamin E, and β -carotene that exist biologically for the regulation of redox homeostasis are also depressed during hyperglycemia induced oxidative stress³⁵. The consequences of chronic oxidative stress are many that include damage to biological macromolecules such as DNA, lipids, proteins, and carbohydrates, disruption in cellular homeostasis, and generation of other ROS. These processes create further damage that result in many pathological processes of clinical interest that includes the diabetic retinopathy²⁵. However, the exact mechanism by which oxidative stress could contribute to the development of diabetic complications still remains to be clarified.

Role of other signalling pathways

Oxidative stress, besides creating a vicious cycle of damage to macromolecules by amplifying the production of more ROS, also activates other metabolic pathways that are detrimental and promote the development of diabetic retinopathy. These include the Advanced Glycation End products (AGE) pathway¹⁵, the polyol pathway⁵⁶, the hexosamine biosynthesis pathway³¹, the de novo synthesis of diacylglycerol (DAG) leading to the overactivation of several isoforms of protein kinase C [Protein Kinase C (PKC) pathway]⁷³¹⁰²⁶⁸¹¹⁰ besides several other that include alteration in the expressions of Vascular Endothelial Growth Factor (VEGF)²¹ and insulin-like growth factor-1 (IGF-1)²⁷, and elevation in mitochondrial overproduction of superoxide and mitochondrial dysfunctions (Fig. 2)⁶⁰.



Fig. 2: The cross-talk between different metabolic pathways involved in the development of diabetic retinopathy. Hyperglycemia induced metabolic dysregulation in different signaling pathways ultimately converge into the production of high levels of ROS, which in conjunction with inflammatory insults lead to the development diabetic retinopathy.

Advanced glycation end products (age) pathway

Advanced Glycation End products (AGE) are a heterogeneous group of compounds that are produced as a result of non-enzymatic condensation reactions between reducing sugars and α -amino groups or N-terminal groups of amino acids, preferentially lysine and arginine¹⁰⁹. The heterogeneity of AGEs comes from the fact that majority of these products come from an array of precursor molecules with variable chemical structure. The intermediates that are mostly involved in the formation of AGEs include dicarbonyls such as 1-, 3-, or 4-deoxyglucosones, glyoxal, and methylglyoxal. These are highly reactive intermediates that are upregulated during hyperglycemia, and react with proteins and form intramolecular or intermolecular AGEs¹⁰⁴¹⁰⁵. The AGEs are also formed on the amino groups of lipids and DNA. AGE formation can lead to the alterations in function, activity, and degradation of both intracellular and extracellular proteins via chemical rearrangement and intramolecular and intermolecular cross-linking. For example, it causes the cross-linking of proteins such as collagen or elastin in the wall of blood vessels thereby altering its structural integrity.

Besides being able to crosslink a number of proteins and alter their physical properties, AGEs interact with a specific receptor, RAGE on vascular cells to activate multiple signaling pathways (e.g. protein kinase C [PKC] and MAP kinase pathways) and activate nuclear factors (e.g. NF- κ B and Cyclic AMP-Responsive Element-Binding protein; CREB, resulting in an increase in ROS production and elaboration of inflammatory factors that cause further damage to the cells¹⁵. There is an increased accumulation of AGE and its receptor, RAGE in the retinal microvasculature in diabetes¹⁰³. AGEs are irreversibly formed and accumulate within retinal capillary cells in the late stages of retinopathy. The AGEs also increase oxidative stress in the retinal vascular cells and initiates a sequence of events leading to retinal capillary cell apoptosis via activation of NF- κ B and caspase-3⁶¹. It may also lead to nitration of proteins which can inactivate mitochondrial and cytosolic proteins; disrupt protein assembly and functions, and increase apoptosis, ultimately leading to pathological consequences and cellular damage²³.

Polyol Pathway

The polyol pathway becomes active under hyperglycemic conditions⁴¹⁸⁷. Elevation in glucose activates the enzyme Aldose Reductase (AR), the first and rate-limiting enzyme in the pathway which reduces glucose to sorbitol using NADPH as a cofactor. As NADPH is used, increased aldose reductase activity (polyol pathway) can therefore alter NADPH/NADP balance and decrease the availability of NADPH cofactor for glutathione reductase, which is critical for the maintenance of the intracellular pool of antioxidant reduced glutathione (GSH). This decreases the capability of cells to respond to oxidative stress¹². Sorbitol produced from glucose by AR is then metabolized to fructose by sorbitol dehydrogenase that uses NAD^+ as a cofactor. Since the conversion of sorbitol to fructose requires reduction of NAD^+ to NADH, sorbitol dehydrogenase leads to an increased ratio of NADH/ NAD^+ , a condition termed "pseudohypoxia" that has been linked to a multitude of metabolic and signaling changes known to alter cell function¹¹². The increased amounts of NADH lead to higher levels of oxidized triose phosphates, precursors of AGEs and of diacylglycerol (DAG) through α -glycerol-3-phosphate. Thus, activities through the AGE and PKC pathways are enhanced²⁶. The excess NADH may become a substrate for NADH oxidase, and this would lead to the generation of intracellular oxidant species⁷². Sorbitol is a polyhydroxylated, and strongly hydrophilic alcohol that does not diffuse readily through cell membranes and therefore accumulates intracellularly with possible osmotic consequences³⁰.

Hexosamine Pathway

Another pathway that plays an important role in the development of hyperglycemia induced diabetic complications including the diabetic retinopathy is the hexosamine pathway⁹⁹⁵⁸. This pathway is yet another outcome of hyperglycemia and increased superoxide levels that are characteristic of diabetes¹⁴. The inhibition of glyceraldehyde-3-phosphate dehydrogenase (GAPDH) and hyperglycemia-induced mitochondrial superoxide overproduction plays an important role in the activation of the hexosamine pathway, presumably by diverting the upstream metabolite fructose-6-phosphate from glycolysis to glucosamine formation⁷¹. Inhibition of GAPDH can result in increased levels of glycolytic metabolite, glyceraldehyde-3-phosphate that can activate the AGE pathway by activating intracellular AGE precursor methylglyoxal⁷⁸¹¹⁰. This pathway results in the production of UDP-N-acetyl glucosamine which is a substrate used for the post-translational modification of intracellular factors including transcription factors³¹. In addition, GAPDH can also be modified by direct glycation and by nitration⁵⁰¹⁴ and over-expression of manganese SOD (MnSOD) decreases the activation of GAPDH.

Protein kinase C (PKC) pathway

The Protein Kinase C (PKC) pathway also plays an important role in the pathogenesis of diabetic retinopathy⁴⁹¹⁰⁹⁶⁰. Hyperglycemia can activate the PKC pathway directly via diacylglycerol (DAG) synthesis and indirectly through production of ROS or activation of the AGE/RAGE pathway and polyol pathway products⁹¹¹⁰²¹⁰¹⁶²⁹⁴. PKC has many isoforms and all the classic and novel PKC isoforms are activated by DAG, but primarily the α and δ isoforms appear to be involved in diabetes⁶⁷.

Activated PKC, particularly PKC β can bring about a variety of changes characteristic of diabetic retinopathy that include increasing vessel permeability, blood flow by virtue of its ability to alter the balance of vasodilatory/ vasoconstrictive factors [e.g. endothelial nitric oxide synthase(eNOS), endothelin-1(ET-1)], alteration of hormone and growth factor receptor recycling, stimulation of neovascularization, induction of adhesion molecule expression [e.g. vascular cell adhesion molecule-1(VCAM-1)] endothelial proliferation and apoptosis, and regulating the action of several factors such as VEGF, IGF-1, Connective Tissue Growth Factor (CTGF) and transforming growth factor β 1(TGF β 1)⁸⁹¹¹⁴¹¹⁰¹⁰¹⁹⁴. De novo synthesis of DAG comes largely from glycolytic intermediates and stepwise acylation of glycerol-3-phosphate. DAG can also be synthesized through the phospholipase pathways activated by growth factors, cytokines, and hormones, such as angiotensin II, which are elevated in diabetes⁷⁷¹¹¹¹¹⁹.

Several of these pathways may also mediate oxidative stress creating an interrelated connection with oxidative stress as well as with other pathways that amplify tissue damage even further. However, it is difficult to pinpoint which pathway/s is/are critical to the development of diabetic retinopathy. More likely, no single metabolic dysfunction seems to be the sole contributor and possibly all pathways interact to create the histopathology changes seen in diabetic retinopathy.

Inflammation in Diabetic Retinopathy

The inflammatory nature of Diabetic Retinopathy (DR) was first suggested by the 1960s finding that there is a lower incidence of DR among diabetic patients taking salicylates for rheumatoid arthritis⁹³. This has been complemented by the subsequent work that has provided further support for this linkage and has identified many characteristic inflammatory features that accompany the progress of DR in patients and in animal models.

The Inflammatory features that characterize diabetic retinopathy include increased blood flow and vascular permeability, tissue (macular) edema, neovascularization⁹, increased expression of inflammatory mediators⁴⁵⁷⁴³⁹⁴⁰¹¹⁶¹⁸²²⁷⁹⁹²¹⁰⁷¹⁰⁸²⁹⁴⁶, accelerated retinal neural¹¹ and microvascular cell death, macrophage infiltration³³²², microglial cell activation⁹⁸¹¹⁷⁶⁹, increased leukocyte adhesion⁸¹⁵², complement activation¹¹⁸, Fas ligand upregulation⁵³ and acute-phase response protein expression⁴³.

The inflammatory response in early diabetes includes cytokine release, which activates key transcriptional regulators, including nuclear factor kappa B (NF- β). NF- β is a widely expressed inducible transcription factor that is an important regulator of many genes involved in mammalian inflammatory and immune responses, proliferation and apoptosis. It regulates expression of many proinflammatory molecules at the level of gene transcription including inducible Nitric Oxide Synthase (iNOS), cyclooxygenase 2 (COX-2), interleukin-1 (IL-1), Tumor Necrosis Factor alpha (TNF- α) and intracellular adhesion molecule (ICAM). The important role of NF- β in the pathogenesis of early stages of diabetic retinopathy is well supported by the fact that the inhibition of proteins whose expression is regulated by NF- β (such as iNOS and ICAM as already mentioned) inhibit diabetes induced degeneration of retinal capillaries. Also, it is now well established that the compounds known to inhibit NF- β likewise inhibit the development of the retinopathy. For example, several different antioxidants which inhibit the development of capillary degeneration and pericyte loss in retinas of diabetic rats⁶³ also inhibit the diabetes-induced activation of retinal NF- β ⁹⁶. Likewise, low-intermediate doses of salicylates (aspirin, sodium salicylate, and sulfasalazine) which inhibited NF- β activation in retinas of diabetic rats, also inhibited expression of inflammatory mediators like iNOS and ICAM-1, and capillary degeneration and pericyte loss in those animals⁶³⁶⁶. Aspirin is known to inhibit the production of prostaglandins also, but salicylate and sulfasalazine have much less of this activity, suggesting that the common action of these three salicylates to inhibit retinopathy in diabetes was primarily not mediated by inhibition of prostaglandins. The proinflammatory molecules such as inducible nitric oxide synthase (iNOS), cyclooxygenase 2 (COX-2), interleukin-1 (IL-1), tumor necrosis factor alpha (TNF- α) and Intracellular Adhesion Molecule (ICAM) whose expression is regulated by NF- β contribute in their own capacity towards the development of diabetic retinopathy. The increased expression of cell adhesion molecules, such as ICAM-1, for example, promotes the attraction and adhesion of leukocytes to the retinal vascular wall, a process known as leukostasis.

Leukostasis is linked to increased vascular permeability possibly through the mechanisms that may involve direct action on tight junction disruption²⁸¹⁷ as well as local upregulation of Vascular Endothelial Growth Factor (VEGF), either from the hypoxia induced by nonperfusion or from release by the leukocytes themselves⁴⁸³⁸⁴²⁴⁷. However, it is the concomitant increase in endothelial cell injury and subsequent cell death that is mainly responsible for the process⁵¹⁵²⁵³.

Advances & hurdles in the treatment of DR

Recent advances in elucidating the molecular mechanisms of disease have led to the development of an unprecedented number of potentially disease-modifying pharmacologic agents for DR. To a greater or lesser extent, all of the proposed treatments for DR inhibit processes that increase fluid leakage, cellular proliferation, or inflammation independently of hyperglycemic control. Treatments currently proposed for DR and their associated effects are grouped into 7 drug classes: antioxidants, Aldose Reductase Inhibitors (ARIs), growth hormone inhibitors, renin-angiotensinaldosterone system inhibitors, steroids, Vascular Endothelial Growth Factor (VEGF) inhibitors, and Protein Kinase C (PKC) inhibitors. Ideally, it is desirable for a drug to have a highly specific effect on abnormal physiological processes while leaving normal physiological processes unimpaired. However, numerous cytokines, growth factors, and other substances contribute to each aspect of retinal tissue damage in DR and they come from different metabolic pathways. Furthermore, some of these pathways and factors are essential to normal physiologic functioning. Thus, suppressing abnormal molecular activity but still permitting normal physiologic

function is a significant challenge in designing disease-modifying drugs for DR.

References

1. The effect of intensive treatment of diabetes on the development and progression of long-term complications in insulin-dependent diabetes mellitus. The Diabetes Control and Complications Trial Research Group. *N Engl J Med.* 1993, 329:977-986.
2. Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33). UK Prospective Diabetes Study (UKPDS) Group. *Lancet.* 1998, 352:837-853.
3. Report of the expert committee on the diagnosis and classification of diabetes mellitus. *Diabetes Care.* 2003, 26 Suppl 1:S5-20.
4. Adamis AP, Miller JW, Bernal MT, D'Amico DJ, Folkman J, Yeo TK and Yeo KT. 1994. Increased vascular endothelial growth factor levels in the vitreous of eyes with proliferative diabetic retinopathy. *Am J Ophthalmol.*, 118:445-450.
5. Aiello LP, Avery RL, Arrigg PG, Keyt BA, Jampel HD, Shah ST, Pasquale LR, Thieme H, Iwamoto MA, Park JE. 1994. Vascular endothelial growth factor in ocular fluid of patients with diabetic retinopathy and other retinal disorders. *N Engl J Med.*, 331:1480-1487.
6. Aiello LP, Gardner TW, King GL, Blankenship G, Cavallerano JD, Ferris FL, 3rd and Klein R. 1998. Diabetic retinopathy. *Diabetes Care* 21:143-156.
7. Amos AF, McCarty DJ, Zimmet P. 1997. The rising global burden of diabetes and its complications: estimates and projections to the year 2010. *Diabet Med.*, 14 Suppl 5:S1-85.
8. Anderson RE, Rapp LM, Wiegand RD. 1984. Lipid peroxidation and retinal degeneration. *Curr Eye Res.* 3:223-227.
9. Antonetti DA, Barber AJ, Bronson SK, Freeman WM, Gardner TW, Jefferson LS, Kester M, Kimball SR, Krady JK, LaNoue KF, Norbury CC, Quinn PG, Sandirasegarane L and Simpson IA. 2006. Diabetic retinopathy: seeing beyond glucose-induced microvascular disease. *Diabetes* 55:2401-2411.
10. Aylward GW. 2005. Progressive changes in diabetics and their management. *Eye* 19:1115-1118.
11. Barber AJ, Lieth E, Khin SA, Antonetti DA, Buchanan AG and Gardner TW. 1998. Neural apoptosis in the retina during experimental and human diabetes. Early onset and effect of insulin. *J Clin Invest.*, 102:783-791.
12. Barnett PA, Gonzalez RG, Chylack LT, Jr. and Cheng HM. 1986. The effect of oxidation on sorbitol pathway kinetics. *Diabetes* 35:426-432.
13. Baynes JW, Thorpe SR. 1999. Role of oxidative stress in diabetic complications: a new perspective on an old paradigm. *Diabetes* 48:1-9.
14. Beisswenger PJ, Drummond KS, Nelson RG, Howell SK, Szwegold BS and Mauer M. 2005. Susceptibility to diabetic nephropathy is related to dicarbonyl and oxidative stress. *Diabetes* 54:3274-3281.
15. Beisswenger PJ, Howell SK, Smith K and Szwegold BS. 2003. Glyceraldehyde-3-phosphate dehydrogenase activity as an independent modifier of methylglyoxal levels in diabetes. *Biochim Biophys Acta.* 1637:98-106.
16. Biessels GJ, van der Heide LP, Kamal A, Bleys RL and Gispen WH. 2002. Ageing and diabetes: implications for brain function. *Eur J Pharmacol.* 441:1-14.
17. Bolton SJ, Anthony DC, Perry VH. 1998. Loss of the tight junction proteins occludin and zonula occludens-1 from cerebral vascular endothelium during neutrophil-induced blood-brain barrier breakdown in vivo. *Neuroscience* 86:1245-1257.
18. Brooks HL, Jr., Caballero S, Jr., Newell CK, Steinmetz RL, Watson D, Segal MS, Harrison JK, Scott EW and Grant MB. 2004. Vitreous levels of vascular endothelial growth factor and stromal-derived factor 1 in patients with diabetic retinopathy and cystoid macular edema before and after intraocular injection of triamcinolone. *Arch Ophthalmol.*, 122:1801-1807.
19. Brownlee M. 2001. Biochemistry and molecular cell biology of diabetic complications. *Nature* 414:813-820.
20. Brownlee M. 2005. The pathobiology of diabetic complications: a unifying mechanism. *Diabetes* 54:1615-1625.
21. Caldwell RB, Bartoli M, Behzadian MA, El-Remessy AE, Al-Shabrawey M, Platt DH, Liou GI and Caldwell RW. 2005. Vascular endothelial growth factor and diabetic retinopathy: role of oxidative stress. *Curr Drug Targets* 6:511-524.
22. Canataroglu H, Varinli I, Ozcan AA, Canataroglu A, Doran F and Varinli S. 2005. Interleukin (IL)-6, interleukin (IL)-8 levels

and cellular composition of the vitreous humor in proliferative diabetic retinopathy, proliferative vitreoretinopathy, and traumatic proliferative vitreoretinopathy. *Ocul Immunol Inflamm.*, 13:375-381.

23. Cowell RM, Russell JW. 2004. Nitrosative injury and antioxidant therapy in the management of diabetic neuropathy. *J Investig Med.*,52:33-44.
24. Cunha-Vaz J, Faria de Abreu JR, Campos AJ. 1975. Early breakdown of the blood-retinal barrier in diabetes. *Br J Ophthalmol.* 59:649-656.
25. Cutler RG. 2005. Oxidative stress profiling: part I. Its potential importance in the optimization of human health. *Ann N Y Acad Sci.* 1055:93-135.
26. Danis RP, Davis MD. 2008. Proliferative Diabetic Retinopathy. In: Duh E ed., *Contemporary Diabetes: Diabetic Retinopathy* Totowa, New Jersey: Humana Press 29 - 65.
27. DeBosch BJ, Baur E, Deo BK, Hiraoka M and Kumagai AK. 2001. Effects of insulin-like growth factor-1 on retinal endothelial cell glucose transport and proliferation. *J Neurochem.*, 77:1157-1167.
28. Del Maschio A, Zanetti A, Corada M, Rival Y, Ruco L, Lampugnani MG and Dejana E. 1996. Polymorphonuclear leukocyte adhesion triggers the disorganization of endothelial cell-to-cell adherens junctions. *J Cell Biol.* 135:497-510.
29. Demircan N, Safran BG, Soylu M, Ozcan AA and Sizmaz S. 2006. Determination of vitreous interleukin-1 (IL-1) and tumour necrosis factor (TNF) levels in proliferative diabetic retinopathy. *Eye.* 20:1366-1369.
30. Du X, Matsumura T, Edelstein D, Rossetti L, Zsengeller Z, Szabo C and Brownlee M. 2003. Inhibition of GAPDH activity by poly(ADP-ribose) polymerase activates three major pathways of hyperglycemic damage in endothelial cells. *J Clin Invest.* 112:1049-1057.
31. Du XL, Edelstein D, Rossetti L, Fantus IG, Goldberg H, Ziyadeh F, Wu J and Brownlee M. 2000. Hyperglycemia-induced mitochondrial superoxide overproduction activates the hexosamine pathway and induces plasminogen activator inhibitor-1 expression by increasing Sp1 glycosylation. *Proc Natl Acad Sci USA.* 97:12222-12226.
32. Engerman RL, Kern TS. 1995. Retinopathy in animal models of diabetes. *Diabetes Metab Rev.*, 11:109-120.
33. Esser P, Heimann K, Wiedemann P. 1993. Macrophages in proliferative vitreoretinopathy and proliferative diabetic retinopathy: differentiation of subpopulations. *Br J Ophthalmol.*, 77:731-733.
34. Feldman EL. 2003. Oxidative stress and diabetic neuropathy: a new understanding of an old problem. *J Clin Invest.*, 111:431-433.
35. Ford ES, Mokdad AH, Giles WH and Brown DW. 2003. The metabolic syndrome and antioxidant concentrations: findings from the Third National Health and Nutrition Examination Survey. *Diabetes* 52:2346-2352.
36. Frank RN. 2004. Diabetic retinopathy. *N Engl J Med.* , 350:48-58.
37. Frank RN. 1991. On the pathogenesis of diabetic retinopathy. A 1990 update. *Ophthalmology* 98:586-593.
38. Freeman MR, Schneck FX, Gagnon ML, Corless C, Soker S, Niknejad K, Peoples GE and Klagsbrun M. 1995. Peripheral blood T lymphocytes and lymphocytes infiltrating human cancers express vascular endothelial growth factor: a potential role for T cells in angiogenesis. *Cancer Res.*,55:4140-4145.
39. Funatsu H, Yamashita H, Ikeda T, Nakanishi Y, Kitano S and Hori S. 2002. Angiotensin II and vascular endothelial growth factor in the vitreous fluid of patients with diabetic macular edema and other retinal disorders. *Am J Ophthalmol.*, 133:537-543.
40. Funatsu H, Yamashita H, Shimizu E, Kojima R and Hori S. 2001. Relationship between vascular endothelial growth factor and interleukin-6 in diabetic retinopathy. *Retina* 21:469-477.
41. Gabbay KH. 1973. The sorbitol pathway and the complications of diabetes. *N Engl J Med.* 288:831-836.
42. Gaudry M, Bregerie O, Andrieu V, El Benna J, Pocard MA and Hakim J. 1997. Intracellular pool of vascular endothelial growth factor in human neutrophils. *Blood.* 90:4153-4161.
43. Gerhardinger C, Costa MB, Coulombe MC, Toth I, Hoehn T and Grosu P. 2005. Expression of acute-phase response proteins in retinal Muller cells in diabetes. *Invest Ophthalmol Vis Sci.*, 46:349-357.
44. Harhaj NS, Antonetti DA. 2004. Regulation of tight junctions and loss of barrier function in pathophysiology. *Int J Biochem Cell Biol.*, 36:1206-1237.
45. Haskins K, Bradley B, Powers K, Fadok V, Flores S, Ling X, Pugazhenth S, Reusch J and Kench J. 2003. Oxidative stress in type 1 diabetes. *Ann N Y Acad Sci*1005:43-54.

46. Hernandez C, Fonollosa A, Garcia-Ramirez M, Higuera M, Catalan R, Miralles A, Garcia-Arumi J and Simo R. 2006. Erythropoietin is expressed in the human retina and it is highly elevated in the vitreous fluid of patients with diabetic macular edema. *Diabetes Care* 29:2028-2033.
47. Horiuchi T, Weller PF. 1997. Expression of vascular endothelial growth factor by human eosinophils: upregulation by granulocyte macrophage colony-stimulating factor and interleukin-5. *Am J Respir Cell Mol Biol.*, 17:70-77.
48. Iijima K, Yoshikawa N, Connolly DT and Nakamura H. 1993. Human mesangial cells and peripheral blood mononuclear cells produce vascular permeability factor. *Kidney Int.*, 44:959-966.
49. Ishii H, Jirousek MR, Koya D, Takagi C, Xia P, Clermont A, Bursell SE, Kern TS, Ballas LM, Heath WF, Stramm LE, Feener EP and King GL. 1996. Amelioration of vascular dysfunctions in diabetic rats by an oral PKC beta inhibitor. *Science* 272:728-731.
50. Ishii T, Sunami O, Nakajima H, Nishio H, Takeuchi T and Hata F. 1999. Critical role of sulfenic acid formation of thiols in the inactivation of glyceraldehyde-3-phosphate dehydrogenase by nitric oxide. *Biochem Pharmacol.* 58:133-143.
51. Jousseaume AM, Murata T, Tsujikawa A, Kirchhof B, Bursell SE and Adamis AP. 2001. Leukocyte-mediated endothelial cell injury and death in the diabetic retina. *Am J Pathol.* 158:147-152.
52. Jousseaume AM, Poulaki V, Le ML, Koizumi K, Esser C, Janicki H, Schraermeyer U, Kociok N, Fauser S, Kirchhof B, Kern TS and Adamis AP. 2004. A central role for inflammation in the pathogenesis of diabetic retinopathy. *FASEB J.* 18:1450-1452.
53. Jousseaume AM, Poulaki V, Mitsiades N, Cai WY, Suzuma I, Pak J, Ju ST, Rook SL, Esser P, Mitsiades CS, Kirchhof B, Adamis AP and Aiello LP. 2003. Suppression of Fas-FasL-induced endothelial cell apoptosis prevents diabetic blood-retinal barrier breakdown in a model of streptozotocin-induced diabetes. *FASEB J.* 17:76-78.
54. Jousseaume AM, Smyth N, Niessen C. 2007. Pathophysiology of diabetic macular edema. In: Lang GE ed., *Diabetic Retinopathy* Basel: Karger 1-12.
55. Kanwar M, Chan PS, Kern TS and Kowluru RA. 2007. Oxidative damage in the retinal mitochondria of diabetic mice: possible protection by superoxide dismutase. *Invest Ophthalmol Vis Sci.* 48:3805-3811.
56. Kern TS, Kowluru RA, Engerman RL. 1994. Abnormalities of retinal metabolism in diabetes or galactosemia: ATPases and glutathione. *Invest Ophthalmol Vis Sci.* 35:2962-2967.
57. Klein R, Klein BE, Moss SE, Davis MD and DeMets DL. 1984. The Wisconsin epidemiologic study of diabetic retinopathy. III. Prevalence and risk of diabetic retinopathy when age at diagnosis is 30 or more years. *Arch Ophthalmol.*, 102:527-532.
58. Kolm-Litty V, Sauer U, Nerlich A, Lehmann R and Schleicher ED. 1998. High glucose-induced transforming growth factor beta1 production is mediated by the hexosamine pathway in porcine glomerular mesangial cells. *J Clin Invest.*, 101:160-169.
59. Kowluru R, Kern TS, Engerman RL. 1994. Abnormalities of retinal metabolism in diabetes or galactosemia. II. Comparison of gamma-glutamyl transpeptidase in retina and cerebral cortex, and effects of antioxidant therapy. *Curr Eye Res.*, 13:891-896.
60. Kowluru RA. 2005. Diabetic retinopathy: mitochondrial dysfunction and retinal capillary cell death. *Antioxid Redox Signal.*, 7:1581-1587.
61. Kowluru RA. 2005. Effect of advanced glycation end products on accelerated apoptosis of retinal capillary cells under in vitro conditions *Life Sci.*, 76:1051-1060.
62. Kowluru RA. 2003. Effect of reinstatement of good glycemic control on retinal oxidative stress and nitrate stress in diabetic rats. *Diabetes* 52:818-823.
63. Kowluru RA, Engerman RL, Case GL and Kern TS. 2001. Retinal glutamate in diabetes and effect of antioxidants. *Neurochem Int.*, 38:385-390.
64. Kowluru RA, Kern TS, Engerman RL. 1997. Abnormalities of retinal metabolism in diabetes or experimental galactosemia. IV. Antioxidant defense system. *Free Radic Biol Med.* 22:587-592.
65. Kowluru RA, Kern TS, Engerman RL and Armstrong D. 1996. Abnormalities of retinal metabolism in diabetes or experimental galactosemia. III. Effects of antioxidants. *Diabetes.* 45:1233-1237.
66. Kowluru RA, Tang J, Kern TS. . 2001. Abnormalities of retinal metabolism in diabetes and experimental galactosemia. VII. Effect of long-term administration of antioxidants on the development of retinopathy. *Diabetes* 50:1938-1942.
67. Koya D, Jirousek MR, Lin YW, Ishii H, Kuboki K and King GL. 1997. Characterization of protein kinase C beta isoform activation on the gene expression of transforming growth factor-beta, extracellular matrix components, and prostanooids in the

glomeruli of diabetic rats. *J Clin Invest.* 100:115-126.

68. Koya D, King GL. 1998. Protein kinase C activation and the development of diabetic complications. *Diabetes* 47:859-866.
69. Krady JK, Basu A, Allen CM, Xu Y, LaNoue KF, Gardner TW and Levison SW. 2005. Minocycline reduces proinflammatory cytokine expression, microglial activation, and caspase-3 activation in a rodent model of diabetic retinopathy. *Diabetes.* 54:1559-1565.
70. Kristinsson JK, Gottfredsdottir MS, Stefansson E. 1997. Retinal vessel dilatation and elongation precedes diabetic macular oedema. *Br J Ophthalmol.* 81:274-278.
71. Laakso M. 1999. Hyperglycemia and cardiovascular disease in type 2 diabetes. *Diabetes.* 48:937-942.
72. Lassegue B, Clempus RE. 2003. Vascular NAD(P)H oxidases: specific features, expression, and regulation. *Am J Physiol Regul Integr Comp Physiol.* 285:R277-297.
73. Lee TS, MacGregor LC, Fluharty SJ and King GL. 1989. Differential regulation of protein kinase C and (Na,K)-adenosine triphosphatase activities by elevated glucose levels in retinal capillary endothelial cells. *J Clin Invest.* 83:90-94.
74. Limb GA, Chignell AH, Green W, LeRoy F and Dumonde DC. 1996. Distribution of TNF alpha and its reactive vascular adhesion molecules in fibrovascular membranes of proliferative diabetic retinopathy. *Br J Ophthalmol.* 80:168-173.
75. Lin DY, Blumenkranz MS, Brothers RJ and Grosvenor DM. 2002. The sensitivity and specificity of single-field nonmydriatic monochromatic digital fundus photography with remote image interpretation for diabetic retinopathy screening: a comparison with ophthalmoscopy and standardized mydriatic color photography. *Am J Ophthalmol.* 134:204-213.
76. Lorenzi M, Gerhardinger C. 2001. Early cellular and molecular changes induced by diabetes in the retina. *Diabetologia.* 44:791-804.
77. Malhotra A, Kang BP, Cheung S, Opawumi D and Meggs LG. 2001. Angiotensin II promotes glucose-induced activation of cardiac protein kinase C isozymes and phosphorylation of troponin I. *Diabetes.* 50:1918-1926.
78. Meister A. 1988. Glutathione metabolism and its selective modification. *J Biol Chem.* 263:17205-17208.
79. Meleth AD, Agron E, Chan CC, Reed GF, Arora K, Byrnes G, Csaky KG, Ferris FL, 3rd and Chew EY. 2005. Serum inflammatory markers in diabetic retinopathy. *Invest Ophthalmol Vis Sci.* 46:4295-4301.
80. Meyerle CB, Chew EY, III FLF. 2008. Nonproliferative Diabetic Retinopathy. In: Duh E ed., *Contemporary Diabetes: Diabetic Retinopathy* Totowa, New Jersey: Humana Press 3-27.
81. Miyamoto K, Khosrof S, Bursell SE, Rohan R, Murata T, Clermont AC, Aiello LP, Ogura Y and Adamis AP. 1999. Prevention of leukostasis and vascular leakage in streptozotocin-induced diabetic retinopathy via intercellular adhesion molecule-1 inhibition. *Proc Natl Acad Sci U S A.* 96:10836-10841.
82. Mizutani M, Kern TS, Lorenzi M. 1996. Accelerated death of retinal microvascular cells in human and experimental diabetic retinopathy. *J Clin Invest.* 97:2883-2890.
83. Mohamed Q, Gillies MC, Wong TY. 2007. Management of diabetic retinopathy: a systematic review. *JAMA.* 298:902-916.
84. Moss SE, Klein R, Klein BE. 1988. The incidence of vision loss in a diabetic population. *Ophthalmology* 95:1340-1348.
85. Mustata GT, Rosca M, Biemel KM, Reihl O, Smith MA, Viswanathan A, Strauch C, Du Y, Tang J, Kern TS, Lederer MO, Brownlee M, Weiss MF and Monnier VM. 2005. Paradoxical effects of green tea (*Camellia sinensis*) and antioxidant vitamins in diabetic rats: improved retinopathy and renal mitochondrial defects but deterioration of collagen matrix glycoxidation and cross-linking. *Diabetes.* 54:517-526.
86. Nishikawa T, Edelstein D, Brownlee M. 2000. The missing link: a single unifying mechanism for diabetic complications. *Kidney Int Suppl.* 77:S26-30.
87. Oates PJ. 2002. Polyol pathway and diabetic peripheral neuropathy. *Int Rev Neurobiol.* 50:325-392.
88. Obrosova IG, Julius UA. 2005. Role for poly(ADP-ribose) polymerase activation in diabetic nephropathy, neuropathy and retinopathy. *Curr Vasc Pharmacol.* 3:267-283.
89. Oikawa T, Shimamura M, Ashino H, Nakamura O, Kanayasu T, Morita I and Murota S. 1992. Inhibition of angiogenesis by staurosporine, a potent protein kinase inhibitor. *J Antibiot (Tokyo).* 45:1155-1160.
90. Olson JA, Strachan FM, Hipwell JH, Goatman KA, McHardy KC, Forrester JV and Sharp PF. 2003. A comparative evaluation of digital imaging, retinal photography and optometrist examination in screening for diabetic retinopathy. *Diabet Med.* 20:528-534.

91. Palumbo EJ, Sweatt JD, Chen SJ and Klann E. 1992. Oxidation-induced persistent activation of protein kinase C in hippocampal homogenates. *Biochem Biophys Res Commun.* 187:1439-1445.
92. Patel JI, Hykin PG, Gregor ZJ, Boulton M and Cree IA. 2005. Angiopoietin concentrations in diabetic retinopathy. *Br J Ophthalmol.* 89:480-483.
93. Powell ED, Field RA. 1964. Diabetic Retinopathy and Rheumatoid Arthritis. *Lancet.* 2:17-18.
94. Rask-Madsen C, King GL. 2005. Proatherosclerotic mechanisms involving protein kinase C in diabetes and insulin resistance. *Arterioscler Thromb Vasc Biol.* 25:487-496.
95. Robison WG, Jr., Kador PF, Kinoshita JH. 1985. Early retinal microangiopathy: prevention with aldose reductase inhibitors. *Diabet Med.* 2:196-199.
96. Romeo G, Liu WH, Asnaghi V, Kern TS and Lorenzi M. 2002. Activation of nuclear factor-kappaB induced by diabetes and high glucose regulates a proapoptotic program in retinal pericytes. *Diabetes* 51:2241-2248.
97. Rosen P, Nawroth PP, King G, Moller W, Tritschler HJ and Packer L. 2001. The role of oxidative stress in the onset and progression of diabetes and its complications: a summary of a Congress Series sponsored by UNESCO-MCBN, the American Diabetes Association and the German Diabetes Society. *Diabetes Metab Res Rev.*, 17:189-212.
98. Rungger-Brandle E, Dosso AA, Leuenberger PM. 2000. Glial reactivity, an early feature of diabetic retinopathy. *Invest Ophthalmol Vis Sci.* 41:1971-1980.
99. Sayeski PP, Kudlow JE. 1996. Glucose metabolism to glucosamine is necessary for glucose stimulation of transforming growth factor-alpha gene transcription. *J Biol Chem.* 271:15237-15243.
100. Sharma S, Oliver-Fernandez A, Liu W, Buchholz P and Walt J. 2005. The impact of diabetic retinopathy on health-related quality of life. *Curr Opin Ophthalmol.*, 16:155-159.
101. Sheetz MJ, King GL. 2002. Molecular understanding of hyperglycemia's adverse effects for diabetic complications. *JAMA.* 288:2579-2588.
102. Stauble B, Boscoboinik D, Tasinato A and Azzi A. 1994. Modulation of activator protein-1 (AP-1) transcription factor and protein kinase C by hydrogen peroxide and D-alpha-tocopherol in vascular smooth muscle cells. *Eur J Biochem.* 226:393-402.
103. Stitt AW. 2003. The role of advanced glycation in the pathogenesis of diabetic retinopathy. *Exp Mol Pathol.*, 75:95-108.
104. Thornalley PJ. . 1990. The glyoxalase system: new developments towards functional characterization of a metabolic pathway fundamental to biological life. *Biochem J*269: 1-11.
105. Thornalley PJ, Langborg A, Minhas HS. 1999. Formation of glyoxal, methylglyoxal and 3-deoxyglucosone in the glycation of proteins by glucose. *Biochem J.* 344 Pt 1:109-116.
106. Trautner C, Haastert B, Giani G and Berger M. 2001. Incidence of blindness in southern Germany between 1990 and 1998. *Diabetologia* 44:147-150.
107. Watanabe D, Suzuma K, Matsui S, Kurimoto M, Kiryu J, Kita M, Suzuma I, Ohashi H, Ojima T, Murakami T, Kobayashi T, Masuda S, Nagao M, Yoshimura N and Takagi H. . 2005. Erythropoietin as a retinal angiogenic factor in proliferative diabetic retinopathy. *N Engl J Med*353:782-792.
108. Watanabe D, Suzuma K, Suzuma I, Ohashi H, Ojima T, Kurimoto M, Murakami T, Kimura T and Takagi H. 2005. Vitreous levels of angiopoietin 2 and vascular endothelial growth factor in patients with proliferative diabetic retinopathy. *Am J Ophthalmol.*, 139:476-481.
109. Wautier JL, Schmidt AM. 2004. Protein glycation: a firm link to endothelial cell dysfunction. *Circ Res.*, 95:233-238.
110. Way KJ, Katai N, King GL. 2001. Protein kinase C and the development of diabetic vascular complications. *Diabet Med.*, 18:945-959.
111. Wilkinson-Berka JL. 2006. Angiotensin and diabetic retinopathy. *Int J Biochem Cell Biol.*, 38:752-765.
112. Williamson JR, Chang K, Frangos M, Hasan KS, Ido Y, Kawamura T, Nyengaard JR, van den Enden M, Kilo C and Tilton RG. 1993. Hyperglycemic pseudohypoxia and diabetic complications. *Diabetes* 42: 801-813.
113. Wohaieb SA, Godin DV. 1987. Alterations in free radical tissue-defense mechanisms in streptozocin-induced diabetes in rat. Effects of insulin treatment. *Diabetes* 36:1014-1018.
114. Xia P, Aiello LP, Ishii H, Jiang ZY, Park DJ, Robinson GS, Takagi H, Newsome WP, Jirousek MR and King GL. 1996.

Characterization of vascular endothelial growth factor's effect on the activation of protein kinase C, its isoforms, and endothelial cell growth. *J Clin Invest* 98:2018-2026.

115. Yan SF, Ramasamy R, Naka Y and Schmidt AM. 2003. Glycation, inflammation, and RAGE: a scaffold for the macrovascular complications of diabetes and beyond. *Circ Res.*, 93:1159-1169.

116. Yuuki T, Kanda T, Kimura Y, Kotajima N, Tamura J, Kobayashi I and Kishi S. 2001. Inflammatory cytokines in vitreous fluid and serum of patients with diabetic vitreoretinopathy. *J Diabetes Complications* 15:257-259.

117. Zeng XX, Ng YK, Ling EA. 2000. Neuronal and microglial response in the retina of streptozotocin-induced diabetic rats. *Vis Neurosci.*, 17:463-471.

118. Zhang J, Gerhardinger C, Lorenzi M. 2002. Early complement activation and decreased levels of glycosylphosphatidylinositol-anchored complement inhibitors in human and experimental diabetic retinopathy. *Diabetes.* 51:3499-3504.

119. Zhang L, Ma J, Gu Y and Lin S. 2006. Effects of blocking the renin-angiotensin system on expression and translocation of protein kinase C isoforms in the kidney of diabetic rats. *Nephron Exp Nephrol.*, 104:e103-111.