

# Nutrients and Antinutrients Analysis of Albizia lebbeck Seed

December 15, 2010 · Volume 1; Issue 3

Muhammad N O, Jimoh FO, Nafiu MO, Oloyede OB, Salawu MO

N O M, FO J, MO N, OB O, MO S. Nutrients and Antinutrients Analysis of Albizia lebbeck Seed. Bioresearch Bulletin. 2010 Dec 15 [last modified: 2013 Oct 16]. Edition 1.

## Abstract

**Background:** The nutritional and antinutritional contents of Albizia lebbeck seed were investigated. Levels of nutrients like the crude protein, ether extract, crude fibre, ash, mineral and antinutrients like phytate, cyanide, oxalate, saponin, and tannins were determined.

**Results:** The results indicated a high protein level of  $27.30 \pm 0.001$ , ether extract of  $7.50 \pm 0.10$ , crude fibre of  $38.50 \pm 0.01$ , ash content of  $4.20 \pm 0.10$  and NFE (carbohydrate by difference) content of  $19.40 \pm 0.00$ . Minerals like magnesium,  $7.47 \pm 0.07$  ppm, Fe,  $2.80 \pm 0.002$  ppm, Cu,  $1.83 \pm 0.001$  ppm and Se,  $4.886 \pm 1.03$  ppm. The antinutrients discovered include phytate,  $2.91$  mol/kg, cyanide,  $0.338$  mg/kg, oxalate,  $0.0012$  mg/100g and saponin,  $90.00$  mg/100g. However, the presence of tannins was not detected.

**Conclusion:** It is considered that Albizia lebbeck could be a very good source of protein and minerals in animal feeds, if well processed, to reduce or eliminate the antinutritional factors. The high saponin content also shows that it could be a potent aphrodisiac

## Introduction

The search for protein-rich seed is an ongoing process. This is because protein is limiting in the world nutrients today, especially in Africa where people can really afford the animal protein, which of course is superior to plant protein. Even the plant proteins are becoming more expensive because of the competition between man and his animals for these protein sources. Therefore, there is the need to continually search for protein, especially among the lesser known plants, among which is *Albizia lebbeck*.

*Albizia lebbeck* is (L) Benth (Mimosaceae) belongs to the family Fabaceae. It is a medium to large tree, of multistemmed widely spreading habit (about 30 m in diameter) when grown in the open but capable of good log form in plantation. The seeds are brown, flattened,  $7 \times 1.5$  mm and are 3 – 12 seeds per pod. The seeds are freely produced and are relatively large (7,000 – 8,000 seeds/kg)<sup>8</sup>. It is indigenous to the Indian subcontinent and those areas of Southeast Asia with a marked dry season, and to the monsoon areas of Southern Australia<sup>22</sup>.

Although, geographically widespread, it appears to have potential for increasing pastoral production in extensive systems in the wet-dry tropics where the major problem is low feed quality of the basal diet. It has various names in various regions, for instance it is called 'siris', 'koko', 'vagai' (India), 'tekik' (Japanese), 'kikote', 'tarisi' (Sudanese), 'khago', 'kase' (Thai) and East Indian walnut<sup>14</sup>. The nutritive values of the leaf, flower and pod of *A. lebbeck* in rabbit and sheep have been reported<sup>6,14,18,24,26</sup>. However, information on the nutritional values and the antinutritional content of the *A. lebbeck* seed is still very scanty in literature.

## Materials and Methods

### Materials

#### Sources of Materials

Dry pods of *Albizia lebbeck* were picked from the premises of the main campus of the University of Ilorin, Nigeria and authenticated at the Department of Plant Biology, University of Ilorin, Nigeria. The sodium hydroxide, sodium sulphate used were products of East Anglia Chemicals, Hadleigh Ipswich Suffolk; the petroleum ether (60 – 80°C) used was a product of Eagle Scientific Limited, Unit 5, Eldson Rd., Beeston, Nottingham; the chloroform, methanol, ethanol used were products of BDH Chemicals Ltd., Poole, England. Cupric sulphate, 5-hydrate was a product of J.T. Baker Chemicals Co., Phillipsmore, N.Z. All other reagents used were of analytical grade and prepared in the Biochemistry laboratory, University of Ilorin, Nigeria, with all-glass apparatus.

### Methods

#### Processing of Materials

The seeds were removed from their pods, winnowed and ground to powder form using electric grinding machine. The milled seed was subsequently used for analyses.

### Proximate Analysis

The crude protein, ether extract, crude fibre and ash contents of the *A. lebbbeck* seed were determined as described by AOAC<sup>2</sup>.

### Antinutrient Determination

The determination of the phytate, tannin, cyanide and oxalate contents of the milled *A. lebbbeck* seed were carried out using the methods of Wheeler and Ferre<sup>28</sup>, <sup>21517</sup> respectively.

### Mineral Determination

The determination of the levels of inorganic minerals of the milled *A. lebbbeck* seed was carried out using the perchloric acid digestion (wet oxidation) procedure. P and Fe were determined using the colorimetry method while determination of the Zn, Ca, Mn and Mg contents of the seed were carried out using the atomic absorption spectrophotometer<sup>21323</sup>.

## Results

The results of the proximate analysis of the *A. lebbbeck* seed is shown in Table 1. The seed has a high protein and crude fibre contents of 27.3 and 38.5% respectively. Table 2 shows the results of the mineral analysis of *A. lebbbeck* seed. The seed is rich in Mg, Fe and Se, however, a little amount of Cd, a heavy metal, was observed. The antinutrient composition of the *A. lebbbeck* seed is shown in Table 3. The seed was found to contain a high level of saponin and considerable amount of phytate and cyanide.

Nutrient	Percentage Composition
Moisture	3.10 ± 0.001
Crude Protein	27.30 ± 0.001
Ether extract	7.50 ± 0.10
Crude Fibre	38.50 ± 0.01
Carbohydrate (by difference)	19.40 ± 0.00

Table 1. Proximate Composition (% dry matter) of *Albizia lebbbeck* Seed.

Mineral	Concentration (ppm)
Mg	7.47 ± 0.07
Mn	0.31 ± 0.001
Cu	1.83 ± 0.001
Fe	2.80 ± 0.002
Zn	-
Se	4.87 ± 1.03
Cd	0.15 ± 0.001

Table 2. Mineral Composition (ppm) of *Albizia lebbbeck* Seed

Antinutrient	Concentration
Phytate (mol/kg)	2.91 ± 0.003
Cyanide (mg/kg)	0.34 ± 0.001
Oxalate (mg/100g)	0.0012 ± 0.003
Saponin (mg/100g)	90.00 ± 0.40

**Table 3.** Antinutrient Content of *Albizia lebbek* Seed.

## Discussion

Animal feed, today, is becoming costly due to the limitation posed by protein source. This is because the conventional soybeans and groundnut are over-competed for man and his animals. Therefore, the present result showing a high level of protein in the *A. lebbek* seed indicates that it could be harnessed as a source of protein in animal feed. The level of protein in the seed compare favourably well with that of cowpeas (25%)<sup>10</sup>, pigeon pea (20.4%),<sup>16</sup> jack bean (30%)<sup>27</sup>, limabean, bambara groundnut (23 – 26%)<sup>1719</sup>. The crude lipid of (7.5%) in the seed is higher than those of the cowpeas (2.2%)<sup>10</sup>, pigeon pea (1.6%)<sup>16</sup>, jack bean (3.4%)<sup>27</sup>, limabean (29.3%), bambara groundnut<sup>171921</sup>.

Seeds that are very high in fat content have been reported to retard digestion and normal metabolism, therefore, the seed may have long shelf-life with no tendency for rancidity<sup>39</sup>. The seed could also be a good source of magnesium, iron and selenium in animal diets. The significance of these elements cannot be over emphasized. For instance, magnesium is important in virtually all the steps involved in protein and cholesterol synthesis; iron in the formation of the haem component of the haemoglobin of the red blood cells; while selenium is known to be required by the enzyme glutathione peroxidase that protects the cell membrane against lipid peroxidation.

However, the existence of the antinutrients, which are known to have various deleterious effects, ranging from reduction in feed intake, reduction in bioavailability of minerals to causing death of animals<sup>45122025</sup>, is an indication of the limitation of the use of the seed as protein source in animal feed. The toxic compounds may occur in all parts of the plant, but the seed is normally the most concentrated source, especially, the seed coat<sup>120</sup>. The seed coat would equally account for the high crude fibre observed in the seed. For *A. lebbek* to be harnessed as a source of protein and minerals in animal feeds, it may therefore be necessary that the *A. lebbek* seed be subjected to processing techniques. This will either reduce or eliminate its antinutritional factors. Meanwhile, the high saponin content of the seed may potentiates it as an aphrodisiac.

## References

1. Aletor VA, Aladetimi OO. 1989. Compositional evaluation of some cowpea varieties and some underutilized edible legumes in Nigeria. *Die Nahrung* 33:999-1007.
2. AOAC. 1990. In *Official Methods of Analysis*. Association of Official Analytical Chemists. 15th edition. Washington, D.C. 22-45.
3. Atteh JO. 2002. *Principles and practice of livestock feed manufacturing*. ADLEK printers, Ilorin, Nigeria. 45-77.
4. Butler LG. 1989. Effects of condensed tannins on animal nutrition. In: "Chemistry and significance of condensed tannins" Hemingway and Karchesy. Eds. Plenum Press, New York. 391-402.
5. Butler LG. 1992. Antinutrient effect of condensed and hydrolyzable tannins. *Basic Life Sci.* 59:693-698.
6. Dwatmadji TE, Bird AR, Lowry JB. 1992. Nutritive value of *Albizia lebbek*. *Supplements for Agriculture*, 32:273-278.
7. Ene-Obong HN, Carnovale E. 1992. A comparison of the proximate, mineral and amino acid composition of some known and lesser known legumes in Nigeria. *Food Chem.* 43:169-175.
8. Everist SL. 1986. Use of fodder trees and shrubs. Queensland Department of primary industries, information series Q 185015, Brisbane. 20-34.
9. Ewing WR. *Poultry Nutrition*, 4th Ed. W. Ray Ewing Pub. South Pasadena, California. 195:148-508.
10. Gallup WD, Reder RE. 1943. Sprouted cowpea as a source of protein and vitamins. Oklahoma Agricultural Experiment Station, Stillwater. *Academy of Science.* 33-56
11. Gill LS. 1992. *Ethnomedical uses of plants in Nigeria*. Uniben Press, Benin city, Nigeria.
12. Giner-Chavez BI. 1996. Condensed tannins in tropical forages. Ph.D Thesis, Cornell University, Ithaca, NY, USA. 24-35.
13. Gomori G. 1942. A modification of the colorimetric phosphorous determination for use with the photoelectric colorimeter. *J. Lab. Clin. Med.* 27:955-965.
14. Gupta BS. 1981. Studies of the effect of the molasses feeding on the nutritive value of siris (*Albizia lebbek*) tree leaves. *Indian J. Nutrition and Diet.* 18:144-147.

15. Iwuoha CI, Kalu FA. 1995. Calcium oxalate and physico-chemical properties of cocoyam (*Colocasia esculenta* and *Xanthosoma agittifolium*) tuber flours as affected by processing. *Food Chemistry* 54:61-66.
16. Jambunathan R, Singh U, Subramani V. 1984. Grain quality of sorghom, pearl millet, pigeon pea and chicken pea. International Crops Research Institute for semi-arid Tropics, Patancheru, India. 67-89.
17. Joslyn MA. 1970. *Methods in Food Analysis*. 2nd Edition. Academic Press, London. 845.
18. Lowry JB. 1987. Green leaf fractionation by fruit bats; Is this feeding behavior a unique nutritional strategy for herbivores? *Australian Wildlife Research* 16: 203-206.
19. Olaofe O, Umar YO, Adediran GO. 1993. The effect of nematicides on the nutritive value and functional properties of cowpea seeds (*Vigna unguiculata* L. Walp). *Food Chem.*, 46:337-341.
20. Osagie AU. 1998. Antinutritional factors. In: *Nutritional Quality of Plant Foods*. (A.U. Osagie and O.U. Eka eds). Published by Post Harvest Research Unit, University of Benin, Nigeria. 221-224.
21. Oyenuga VA. 1968. *Nigerian foods and feeding stuffs: Their chemistry and nutritive values*. Ibadan University Press, Nigeria. 36-87.
22. Parrota JA. 1988. In *Early growth and yield of Albizia lebeck at coastal site in Perkin-Elmer Corp (1968)*. Analytical method for Atomic absorption spectrometry. Perkin-Elmer, Corp., Norwalk, Connecticut 76-80.
23. Piper CS. 1944. *Soil and Plant Analysis*. Interscience Publishers, Inc., New York. Puerto Rico. Nitrogen-fixing Research Reports. 6:47-49.
24. Pradhan IP, Dayal R. 1981. Farm forestry in agricultural economy. *Indian Forestry*. 107:665-667.
25. Reed JD. 1995. Nutritional toxicology of tannins and related polyphenols in forage legumes. *J. Anim. Sci.* 73, 1516-1528.
26. Schlink AC, Lowry JB, Gibson DS. 1991. Products from tree legumes. A. lebeck as supplement for sheep in the dry tropics. *Proceedings of Australian Society of Animal Production*. 18:546.
27. Udebibie ABI. 1990. Nutritional evaluation of jackbean (*Canavalia ensiformis*) for the Nigeria poultry industry. *Royal Swidish Academy of Science, AMBIO* 19(8):361.
28. Wheeler EL, Ferrel RE. 1971. A method for phytic acid determination in wheat fractions. *Cereal Chem.*, 48:312-316.